




**Better Safe Than
Sorry:
The Biological Basis
of Fetal Alcohol
Syndrome and other
Alcohol-Related
Birth Defects**

When a mother drinks, her unborn child is exposed to alcohol.



Alcohol-Related Birth Defects Include:

- **Fetal Alcohol Syndrome (FAS)**

which is characterized by

1. reduced IQ
2. low birth weight and height
3. typical facial features

- **Fetal Alcohol Effects (FAE)**

which result from maternal alcohol abuse but are found in the absence of the full-blown syndrome


The facial features of Fetal Alcohol Syndrome are:

- Small eyelid openings (palpebral fissures)
- Short, upturned nose
- Long upper lip (from nose to mouth) with a thin red border and a deficient central groove (philtrum)
- Reduced size of the head (microcephaly)



NORMAL

FAS

An illustration of an iceberg floating in dark blue water. The tip of the iceberg is above the water line, and the much larger, jagged base is submerged. The text 'FAS' is written in yellow above the tip, and 'ARBDS' is written in yellow on the submerged part. To the right of the iceberg, there is a block of yellow text.

FAS

**Full-blown fetal
alcohol syndrome
(FAS)**

ARBDS

**represents only the
“tip of the iceberg”
relative to all
alcohol-related birth
defects (ARBDS).**

**MATERNAL ALCOHOL ABUSE
IS THE LEADING KNOWN
CAUSE OF MENTAL
RETARDATION IN THE
WESTERN WORLD**

Children with alcohol-related birth defects typically have:


attention deficits

- **language difficulties**
- **learning disabilities**
- **impulsive behavior**
- **poor judgment**



PRENATAL ALCOHOL EXPOSURE CAN PERMANENTLY DAMAGE THE BRAIN, AFFECTING IMPORTANT STRUCTURES SUCH AS THE CEREBELLUM AND CORPUS CALLOSUM, AS WELL AS SPECIFIC CELL POPULATIONS IN MANY OTHER REGIONS OF THE BRAIN


Whole brain



cerebral hemispheres


cerebellum

Cross-section



corpus callosum

Visualization of the brain of a normal individual (A) and two with FAS (B, C) shows permanent loss of the tissue indicated by the arrows (portions of the corpus callosum)




Normal

FAS

FAS

Images courtesy of DR. S. Mattson

Similarities in mouse and human embryogenesis provide opportunities to study the effects of alcohol on development



The facial features of Fetal Alcohol Syndrome can be seen in both a child and a mouse fetus that were exposed to alcohol during development.

child with FAS



Narrow forehead

Short palpebral fissures

Small nose

Small midface

Long upper lip with deficient philtrum

mouse fetuses



alcohol-exposed




normal




The amount and timing of maternal alcohol use determine the type and extent of resulting birth defects.

Alcohol can cause malformations and brain abnormalities in embryos that are only three to four weeks old




22 day old human embryo (about 2 mm long, the length of the ear on the US dime)

ALCOHOL KILLS SPECIFIC CELLS IN THE DEVELOPING BRAIN




Arrows surround a portion of the brain of a mouse embryo (viewed from the back) that is at a developmental stage corresponding to a 22-23 day human.




Cells killed by alcohol in the brain of a mouse embryo (at a comparable stage of development to that on the left) have taken up a dark blue stain.

CELLS THAT SHOULD FORM MIDLINE STRUCTURES OF THE BRAIN AND FACE ARE KILLED BY ALCOHOL

Developing
brain and
face




Heart



Mouse embryo (viewed from the front) at a stage corresponding to a 22-23 day old human.

A close-up view of an alcohol-exposed mouse embryo shows cells killed by alcohol that have taken up a dark blue stain


MIDLINE STRUCTURES OF THE FACE AND BRAIN ARE DEFICIENT IN ALCOHOL-EXPOSED MOUSE EMBRYOS AND IN INDIVIDUALS WITH FAS



THE FACE OF A CHILD WITH FULL-BLOWN FAS HAS FEATURES THAT CAN BE CAUSED BY DAMAGE TO MIDLINE STRUCTURES.

COMPARISON OF THE FACE (A) AND INTERIOR OF THE BRAIN (B) OF A NORMAL MOUSE EMBRYO AND ONE DAMAGED BY ALCOHOL (C&D) SHOWS THAT THE NOSTRILS ARE ABNORMALLY POSITIONED (C) AND THE BRAIN IS MISSING MIDLINE STRUCTURES (D)


ALCOHOL KILLS SPECIFIC CELLS IN THE DEVELOPING BRAIN



The pattern of cell death varies with the stage of development.



Cells killed by alcohol have taken up dark blue stain



A cut made through the area outlined by arrows provides a view of the inside of the brain of a 10 days mouse embryo

(corresponding to a 28 day human)

**EXPOSURE TO ALCOHOL
DURING DEVELOPMENT
CAN CAUSE DAMAGE TO
ORGANS AND REGIONS
OTHER THAN THE BRAIN**



**This child with FAS has
a scar from a repaired
cleft lip.**

**Cleft lip can also be caused
by genetic or environmental
agents other than alcohol.**



**Alcohol also caused cleft lip
in this mouse.**



By the ninth week of development the human fetus is about 24 mm. long. Damage caused by alcohol to the brain at this time and until birth can result in abnormal brain function.

Excessive alcohol exposure can cause damage during all stages of prenatal development.

- Pre – implantation: first 2 weeks**
- Embryonic: 3-8 weeks after conception**
- Fetal: from week 9 until birth**



Alcohol can
cause
permanent
damage to a
baby before
most women
realize they are
pregnant.



Alcohol – related birth defects


last a lifetime



Alcohol - related birth defects are expensive:

- **Monetarily** – for treatment, care, and lost productivity. Costs are approximately \$1.4 million over a lifetime for each individual with FAS

- **Socially** – relative to delinquency and to emotional drains on involved families.






???

How much is too much

???



How much alcohol is in a drink?



**12 oz beer = 5 oz wine = shot of liquor
in a mixed drink**

Each contains 0.5 oz of alcohol

WARNING



Some drinks contain more than a “serving” (0.5 oz) of alcohol.



**NO ONE KNOWS WHAT A
“SAFE” AMOUNT OF
ALCOHOL CONSUMPTION
DURING PREGNANCY
MAY BE.**

**Health advisories
urge women who
are planning
pregnancy
or are pregnant
not to drink alcohol.**



Despite warnings, frequent drinking among pregnant women appears to be increasing.

Frequent drinking is defined as 7 or more drinks per week or 5 or more drinks on at least one occasion.





**ALCOHOL-RELATED
BIRTH DEFECTS
ARE PREVENTABLE**



**Better Safe
Than Sorry**

Despite warnings, frequent drinking among pregnant women appears to be increasing.

--	--